FIG. 1





FIG. 2



BEST AVAILABLE COPY

FIG. 3




FIG.4







FIG.5
































FIG. 12



















FIG. 17






FIG. 18












FIG. 21






FIG. 22












FIG. 25





BEST AVAILABLE COPY
































BEST AVAILABLE COPY












FIG. 41


NAMATODE SURVIVAL RATIO RELATIVE TO LIFETIME WHEN FEEDING WATER IS CHANGED







30 31 AVERAGE LIFE SPAN IN ANTIOXIDANT-FUNCTIONING WATER: 20.05 DAYS 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 ~ AVERAGE LIFE SPAN IN PURIFIED WATER: 17.75 DAYS വ O 17 CHANGE IN NAMATODA LIFE SPAN WHEN FEEDING WATER IS CHANGED 12 18 13 18 28 24 23 ស 0 GANTIOXIDANT-FUNCTIONING WATER 5 30 25 20 9 2 0 LIFETIME ■PURIFIED WATER DEATH RATE (NUMBER OF NAMATODA)

BEST AVAILABLE COPY




AVAILABLE COPY






BEST AVAILABLE COPY





TEST AVAILABLE COPY

BEST AVAILABLE COPY



BEST AVAILABLE COPY





BEST AVAILABLE COPY

BEST AVAILABLE COPY





FIG. 51



BEST AVAILABLE COPY